



EVALUATION OF THE EFFICACY OF IMMUNOMODULATING THERAPY FOR ACUTE OBSTRUCTIVE BRONCHITIS IN CHILDREN

Sh.M. Ibatova, A.Kh. Ergashev, D. Kh. Mamatkulova

Samarkand State Medical University, Republic of Uzbekistan

ABSTRACT

Among respiratory diseases, acute obstructive bronchitis (AOB) is common and leads to frequent relapses and severe complications. In young children, under the influence of infectious and other factors, various immunological changes are observed, therefore, the search and implementation of methods that have a corrective effect on the immune system in children with acute obstructive bronchitis is relevant. The aim of the study was to substantiate the use of immunomodulatory therapy in children with acute obstructive bronchitis. 65 children with AOB, 35 patients with acute simple bronchitis and 20 healthy children were examined. Children with acute obstructive bronchitis were aged from 6 months. up to 3 years, of which 39 (60%) were boys, 26 (40%) were girls. The number of T-lymphocytes (SD3), T-helpers (SD4), T-suppressors (SD8), as well as B-lymphocytes (SD19) was determined by a modified method (Yu.F. Garib, 1995). Immunological examination was carried out taking into account the nature of therapy: the 1st group of patients received traditional treatment with the inclusion of T-activin, and the 2nd group of children received only traditional treatment. In order to correct immunological parameters, patients with AOB were prescribed T-activin subcutaneously at the rate of 2 µg/kg of body weight daily for 5 days and the sixth injection a week after the injections. The use of T-activin against the background of traditional therapy has a pronounced positive effect, contributes to a more rapid reduction in the symptoms of intoxication and relief of various complications of the disease. Comparative analysis of the immune response indicators against the background of traditional treatment and with the addition of T-activin revealed a significant increase in B-lymphocytes (DM19) 12.9±0.76%, an increase in FAN 57.9±1.34% and normalization of immunoglobulins A, M, G. The use of T-activin in the complex therapy of children with acute obstructive bronchitis increases the effectiveness of treatment, contributes to the normalization of the immune status and prevents the development of relapses of the disease.

KEYWORDS: acute obstructive bronchitis, children, immune status indicators, immunomodulatory therapy.

INTRODUCTION

Among respiratory diseases, acute obstructive bronchitis (AOB) is widespread, leading to frequent relapses and severe complications. Under the influence of an infectious factor and other agents, various immunological changes are observed in children, and the ability to develop full-fledged post-infection immunity also sharply decreases [1,4,5,7,9]. Therefore, a promising area of research is the search and implementation of methods that have a corrective effect on the immune system in children with acute obstructive bronchitis [2,3, 6,8,10].

PURPOSE OF THE STUDY

To substantiate the use of immunomodulatory therapy in children with acute obstructive bronchitis.

MATERIALS AND METHODS

65 children with AOB, 35 patients with acute simple bronchitis and 20 healthy children were examined. Children with acute obstructive bronchitis were aged from 6 months. up to 3 years, of which 39 (60%) were boys, 26 (40%) were girls. The diagnosis of AOB was established according to the classification adopted in 1996 at the Russian Symposium of Pediatric Pulmonologists.



The immunological study was carried out in the clinical laboratory of the SamMI clinic. The number of T-lymphocytes (SD3), T-helpers (SD4), T-suppressors (SD8), as well as B-lymphocytes (SD19) was determined by a modified method (Yu.F. Garib, 1995). The concentration of serum immunoglobulins A, M, G in peripheral blood was determined by the method of Mancini et al (1965). The phagocytic activity of neutrophils was studied using latex particles (Petrov R.V., 1988).

Immunological examination was carried out taking into account the nature of therapy: the 1st group of patients received traditional treatment with the inclusion of T-activin, and the 2nd group of children received only traditional treatment.

RESEARCH RESULTS

It has been established that in most children the disease occurs at the age of 3 months to 1 year. An analysis of family and hereditary history showed that 32% of sick children were born from related marriages, in 46.5% of children, relatives suffered from allergic diseases. An analysis of the initial premorbid background showed that in children with AOB, allergic diathesis was observed in 54.9%, anemia in 81.9%, rickets in 51.0%, paratrophy in 12.5% and malnutrition of I-II degrees – at 48.7%. It was revealed that the average body weight at birth in children with AOB significantly exceeded (more than 3.5 kg) those in children with acute simple bronchitis and the control group.

The main changes in cellular immunity were expressed in a decrease in the number of T-lymphocytes (DM3) 45.2 ± 0.8 compared to children in the control group $57.3 \pm 0.9\%$ ($p < 0.01$). More often there was an increase in the content of B-lymphocytes (DM19) in patients with OOB 18.1 ± 0.3 ($p < 0.01$), which is significantly higher than data with acute bronchitis $16.1 \pm 0.7\%$ ($p < 0.01$) and in the control group ($p < 0.01$). There was a trend towards a decrease in T-suppressors (DM8) in relative and absolute terms in AOB in children.

The phagocytic activity of neutrophils in the acute period of the disease is significantly inhibited in children with AR 45.1 ± 0 ($p < 0.01$). A particularly pronounced decrease in the increase in the phagocytic activity of neutrophils (FAN) was observed in children with relapses (3-4 times a year) of acute obstructive bronchitis. There was also a significant decrease in the phagocytosis index and the indicator of completed phagocytosis. Changes in the humoral link of immunity were accompanied by a decrease in the concentration of IgA ($p < 0.01$) and IgG ($p < 0.01$). An increase in the concentration of IgM ($p < 0.01$) in children with AOB indicates that during the peak of the disease, the immune response is provided mainly by antibodies of the IgM class.

In children with AOB who were on the traditional method of treatment, the improvement in clinical symptoms and immunological parameters was less pronounced. Thus, the level of T-lymphocytes ($p < 0.01$) remained low, the levels of B-lymphocytes ($p < 0.01$) were high. The content of immunoglobulins did not reach the levels of healthy children.

In order to correct immunological parameters, patients with AOB were prescribed T-activin subcutaneously at the rate of 2 $\mu\text{g}/\text{kg}$ of body weight daily for 5 days and the sixth injection a week after the injections. The use of T-activin against the background of traditional therapy has a pronounced positive effect, contributes to a more rapid reduction in the symptoms of intoxication and relief of various complications of the disease. Comparative analysis of the immune response indicators against the background of traditional treatment and with the addition of T-activin revealed a significant increase in B-lymphocytes (DM19) $12.9 \pm 0.76\%$, an increase in FAN $57.9 \pm 1.34\%$ and normalization of immunoglobulins A, M, G.



CONCLUSION

Thus, the use of T-activin in the complex therapy of children with acute obstructive bronchitis increases the effectiveness of treatment, contributes to the normalization of the immune status and prevents the development of relapses of the disease.

BIBLIOGRAPHY

1. Zaitseva O.V. *Broncho-obstructive syndrome in children. / Pediatrics. 2015. No. 4. -p.94-104.*
2. Ibatova Sh.M., Mamatkulova F.Kh. *Broncho-obstructive syndrome in children: prevalence of difficulties in differential diagnosis and prognosis. / Problems of biology and medicine. Samarkand. 2019, Number 3. -p.233-236.*
3. Ibatova Sh. M., Mamatkulova F. Kh., Ruzikulov N.Y. *The Clinical Picture of Acute Obstructive Bronchitis in Children and the Rationale for Immunomodulatory Therapy. International Journal of Current Research and Review. Vol 12 Issue 17. September 2020. - P.152-155.*
4. Ibatova Sh. M., F. Kh. Mamatkulova, N. B. Abdukadirova, Yu. A. Rakhmonov, M. M. Kodirova. *Risk Factors for Development of Broncho-Ostructive Syndrome in Children. International Journal of Current Research and Review. Vol 12. Issue 23 December 2020.-P. 3-6.*
5. Ibatova Sh.M., Mamatkulova F.Kh., Rakhmonov Y.A., Shukurova D.B., Kodirova M.M. *Assessment of the Effectiveness of Treatment of Rachit in Children by Gas-Liquid Chromatography. International Journal of Current Research and Review. Vol 13, Issue 06, 20 March 2021. P.64-66.*
6. Sh.M. Ibatova, F.Kh. Mamatkulova, D.B. Shukurov. *Identification of risk factors for the development of broncho-obstructive syndrome in young children. Journal "Biomedicine and Practice". Tashkent. No. SI-2. 2020. 4- part. Special issue. pp.481-485.*
7. Sh.M. Ibatova, F.Kh. Mamatkulova, N.Y. Ruzikulov, Yu.A. Rakhmonov. *Bronchoo structive syndrome in children: prevalence and difficulties of differential diagnostics. ACADEMICIA: An International Multidisciplinary Research Journal 2021, P. 87-92.*
8. Ibatova Sh. M., Abdurasulov F.P., Mamutova E.S. *Some aspects of diagnostics of out-of-social pneumonia in children indications for hospitalization. EPRA International Journal of Research and Development (IJRD) Volume: 6 | Issue: 4 | April 2021. P. 242-244.*
9. Ibatova Sh.M., Mamatkulova F.Kh., Mukhamadiev N.K. *State of immunity in chronic obstructive pulmonary disease in children. CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES Volume: 02 Issue: 05 | Sep-Oct 2021 ISSN: 2660-4159. P. 103-107.*
10. Ramazanova A.B., Ibatova Sh.M. *Determination of the level of immunoglobulins in blood serum in infants, depending on the nature of feeding. Scientific journal "Biomedicine and practice". Tashkent. No. SI-2. 2020. 5th part. Special issue. pp. 476-480.*
11. Rakhimov S.A., Zakirov U.I. *Indicators of digital dermatoglyphics of children with obstructive bronchitis of the Uzbek pediatrician population. Pediatrics. 2011. №3. - FROM. 22-26.*