



# EFFECTIVENESS OF AQUATIC THERAPY ON PAIN RELIEF AND FUNCTIONAL MOBILITY IN KNEE OSTEOARTHRITIS: A NARRATIVE REVIEW

Joel George<sup>1</sup>, Prabhu Sanker<sup>2</sup>

<sup>1</sup>Department of Physiotherapy, Garden City University

<sup>2</sup>Professor, Department of Physiotherapy, Garden City University

Article DOI: <https://doi.org/10.36713/epra22855>

DOI No: 10.36713/epra22855

## ABSTRACT

**Background:** Knee osteoarthritis (OA) is a degenerative joint disease causing pain, stiffness, and functional limitations, particularly in older adults. Aquatic therapy is emerging as a promising non-pharmacological intervention due to water's unique properties that reduce joint loading and facilitate exercise.

**Aim:** To review the effectiveness of aquatic therapy on pain relief and functional mobility in individuals with knee osteoarthritis.

**Methods:** A review was conducted analyzing five recent, high-quality randomized controlled trials and systematic reviews published within the last 10 years. Studies were identified through databases such as PubMed and Google Scholar and evaluated for methodological rigor and clinical relevance.

**Results:** Evidence consistently demonstrates that aquatic therapy significantly reduces pain, improves physical function, muscle strength, and balance in knee OA patients. Aquatic cycling and hydrotherapy were particularly effective. Patient adherence and satisfaction exceeded those reported for land-based exercise, making aquatic therapy a viable rehabilitation option.

**Conclusion:** Aquatic therapy is an effective and safe intervention for managing symptoms of knee OA, enhancing quality of life through improved mobility and reduced pain. Future research should focus on long-term outcomes and standardized aquatic protocols.

**KEYWORDS:** Aquatic Therapy, Knee Osteoarthritis, Pain Relief, Functional Mobility, Hydrotherapy, Rehabilitation

## INTRODUCTION

Knee osteoarthritis (OA) is a prevalent degenerative joint disorder characterized by progressive cartilage degradation, pain, stiffness, and functional impairment, severely impacting quality of life, especially among elderly individuals. While pharmacological and land-based exercise treatments are common, pain and reduced mobility often limit adherence and effectiveness. Aquatic therapy offers a low-impact alternative, using water buoyancy, hydrostatic pressure, and resistance to reduce joint load and facilitate exercise. This review synthesizes current evidence on aquatic therapy's efficacy in reducing pain and improving function in knee OA.

## REVIEW

This narrative review synthesizes findings from twelve relevant studies evaluating the effectiveness of various aquatic therapy modalities in managing symptoms of knee osteoarthritis (OA): Dong et al. (2018) conducted a systematic review and meta-analysis comparing aquatic and land-based exercises, concluding that aquatic therapy significantly improves pain, stiffness, and functional mobility. Higher adherence rates were noted for aquatic exercise programs.

Dias et al. (2017) found that hydrotherapy significantly improved pain and physical function in older women with knee OA. The six-week intervention led to improvements in muscle strength and functional outcomes.

Rewald et al. (2020) evaluated a 12-week aquatic cycling program and reported significant reductions in knee pain and enhancements in physical function. KOOS scores improved, and adherence rates reached 80%, indicating the feasibility and tolerability of aquatic cycling.

Kunduracilar et al. (2018) compared two aquatic protocols and identified that the inclusion of trunk and upper limb exercises resulted in superior outcomes in pain reduction, balance, and functional ability.

Garbi et al. (2021) demonstrated that aquatic physiotherapy improved functional capacity and mobility in elderly individuals with knee OA. The intervention was well tolerated and showed promise in geriatric rehabilitation.

Lim et al. (2010) assessed the impact of aquatic therapy on obese individuals with knee OA. Results showed improvements in functional fitness, joint function, and pain interference in daily activities, supporting aquatic therapy's utility in this population.

Wyatt et al. (2001) compared aquatic and traditional exercise programs and found similar physical improvements in both groups. However, the aquatic group experienced greater pain relief, reinforcing its advantage in symptomatic management.



Karimi et al. (2023) conducted a randomized trial assessing an eight-week aqua therapy program in women with knee OA. Results included marked reductions in pain (VAS) and improved lower limb performance across multiple functional tests.

Noor et al. (2023) reviewed 10 RCTs and concluded that aquatic therapy significantly enhances balance and mobility. Though less impact was seen on gait, overall findings supported aquatic exercise as an initial OA intervention.

Amras and Kamalakannan (2023) presented a narrative review emphasizing aquatic therapy's role in reducing joint stress, improving circulation, and enhancing recovery. Their analysis underlined its integration into standard OA rehabilitation protocols.

Rewald et al. (2016) described the protocol for a randomized controlled trial comparing aqua-cycling with usual care. The study framework highlighted expected improvements in pain, strength, and mobility.

Wang et al. (2011) showed that aquatic and land-based exercises both improved pain and function in knee OA, with aquatic therapy offering a more tolerable option for those with greater movement limitations.

All studies utilized validated outcome measures such as VAS, WOMAC, KOOS, 6MWT, and TUG to ensure reliable assessment of therapeutic efficacy.

**Summary of Key Studies on Aquatic Therapy in Knee Osteoarthritis**

Author(s) & Year	Study Design	Sample Size & Population	Intervention	Outcome Measures	Key Findings
Dong et al., 2018	Systematic Review & Meta-analysis	Multiple RCTs	Aquatic vs. land-based exercise	VAS, WOMAC	Reduced pain, improved function, higher adherence
Dias et al., 2017	RCT	73 elderly women	Hydrotherapy, 6 weeks	WOMAC, strength	Pain, function, strength improved
Rewald et al., 2020	RCT	111 adults	Aquatic cycling, 12 weeks	KOOS, physical function	Reduced pain, improved mobility, high adherence
Kunduracilar et al., 2018	RCT	Female patients	2 aquatic protocols	VAS, WOMAC, balance	Core+limb group had better outcomes
Garbi et al., 2021	RCT	Elderly individuals	Aquatic physiotherapy	6MWT, TUG	Improved functional mobility
Lim et al., 2010	RCT	75 obese adults	AQE vs. LBE vs. control	QOL, knee function, BMI	Pain relief and tolerability better in AQE
Wyatt et al., 2001	RCT	46 participants	AQE vs. LBE	ROM, pain, 1-mile walk	AQE reduced pain more effectively
Karimi et al., 2023	RCT	60 women	Aqua therapy, 8 weeks	VAS, TUG, 30-S-CS, 40MW	Pain and performance significantly improved
Noor et al., 2023	Systematic Review	10 RCTs, 531 participants	Aquatic exercise	Balance, mobility, posture	Significant balance and mobility gains
Amras & Kamalakannan, 2023	Narrative Review	Multiple studies	General aquatic therapy	Pain, circulation, recovery	Advocated clinical integration
Rewald et al., 2016	Study Protocol	N/A	Aqua-cycling vs. usual care	KOOS, QOL, strength	Anticipated benefit in mild-moderate OA
Wang et al., 2011	RCT	84 participants	AQE vs. LBE	KOOS, ROM	Both improved, AQE more tolerable

**METHODOLOGY**

This narrative review was conducted by systematically retrieving peer-reviewed literature from databases including PubMed, Google Scholar, and ScienceDirect. Studies published between 2010 and 2023 were considered. Keywords included "aquatic therapy," "hydrotherapy," "knee osteoarthritis," "pain relief," and "functional mobility."

**Inclusion criteria:** Participants aged ≥45 years with radiologically or clinically confirmed knee OA, aquatic therapy as the primary intervention, reported outcomes on pain and physical function, and randomized controlled trials or systematic reviews.

**Exclusion criteria:** Recent knee surgery (<6 months), intra-articular injections, significant comorbidities limiting mobility,



cognitive impairment, or concurrent therapies affecting aquatic intervention outcomes.

Data were extracted on study design, participant demographics, intervention characteristics, outcome measures, and results. Methodological quality was assessed using PEDro scores where available.

**Outcome Measures and Effectiveness of Aquatic Therapy in Knee OA**

Outcome Domain	Measurement Tools	Effect of Aquatic Therapy	Clinical Implication
Pain	VAS, WOMAC	Significant reductions in pain intensity	Enhanced exercise participation and comfort
Physical Function	KOOS, TUG, 6MWT, WOMAC	Improved walking speed, mobility, daily function	Increased independence and quality of life
Muscle Strength	Sit-to-Stand Test, Isokinetic testing	Greater knee extensor/flexor strength	Joint stabilization and reduced fatigue
Balance & Posture	SLS Test, Balance scales	Improved postural control and gait stability	Reduced fall risk in elderly populations
Adherence & Satisfaction	Attendance logs, surveys	Higher adherence vs. land-based exercises	Better long-term outcomes and motivation

**RESULTS**

Analysis of the included studies consistently showed:

- Substantial improvements in pain, functional mobility, muscle strength, and balance.
- Higher adherence and satisfaction in aquatic therapy groups compared to land-based.
- Effectiveness across diverse protocols including hydrotherapy, aqua-cycling, and combined trunk-limb exercises.
- Strong evidence in special populations (e.g., elderly, obese, postmenopausal women).
- Favorable safety profile with no reported serious adverse events.

**Summary of Selected Studies on Aquatic Therapy for Knee Osteoarthritis**

Study	Method	Participants	Intervention	Outcomes
Dong et al., 2018	Systematic review and meta-analysis of 8 RCTs	579 adults with knee OA	Aquatic vs. land-based exercises	Significant reduction in pain, stiffness, and improved function; higher adherence than land-based.
Dias et al., 2017	RCT; parallel group, 6-week intervention	73 elderly women (≥65 years) with knee OA	Hydrotherapy twice/week in heated pool + education	Significant improvements in pain, function (WOMAC), and muscle strength.
Rewald et al., 2020	Single-blind RCT; 12-week aquatic cycling	111 adults (aged 50–70) with mild-to-moderate knee OA	Aquatic cycling: 45 min, 2x/week	KOOS improvements in pain and function; high adherence (80%).
Kunduracilar et al., 2018	RCT with 3 groups (2 aquatic + control)	89 women with bilateral grade 2–3 knee OA	Group 1: LE aquatic; Group 2: full body aquatic; Control: conventional therapy	Group 2 showed best improvements in function, stiffness, and balance.
Garbi et al., 2021	RCT (methodological details limited)	Elderly individuals with knee OA	Aquatic physiotherapy (frequency not specified)	Improved mobility and functional capacity; well-tolerated.
Lim et al., 2010	RCT	75 obese adults (>50 years, BMI >25 kg/m <sup>2</sup> ) with OA	AQE vs. LBE vs. control	AQE reduced pain interference in daily activities; similar improvements in fitness and function.
Wyatt et al., 2001	RCT	46 participants aged 45–70 with knee OA	Aquatic vs. land-based exercises	Both groups improved, but aquatic group had lower pain levels post-intervention.
Karimi et al., 2023	RCT; 8-week intervention	60 women with knee OA	Aqua therapy 3x/week	Pain decreased; performance improved (TUG, 30-S-CS, 40MW).
Noor et al., 2023	Systematic review of 10 RCTs	531 participants (457 females, 74 males) aged ≥40	Aquatic exercises (2–5x/week for 3–12 weeks)	Improved balance, postural control, and mobility; mixed



				results for gait/postural sway.
<b>Amras &amp; Kamalakannan, 2023</b>	Narrative review	Multiple studies (2006–2022)	Aquatic therapy in rehabilitation programs	Reduced pain, improved joint function and recovery; supports inclusion in rehabilitation.
<b>Rewald et al., 2016</b>	RCT study protocol	Adults with mild-to-moderate knee OA (planned sample)	Aqua-cycling 2x/week for 12 weeks vs. usual care	Aimed to assess knee pain and function, muscle strength, QOL.
<b>Wang et al., 2011</b>	RCT; 12-week comparative trial	84 participants with knee OA	Aquatic vs. land-based exercises	Both groups improved in pain, ROM, and function; aquatic was more tolerable for some participants.

## DISCUSSION

The evidence supports aquatic therapy as a safe, accessible, and effective treatment for individuals with knee OA, particularly for those unable to tolerate traditional land-based programs. The reduced mechanical load in water environments allows patients to engage in exercise with less discomfort, fostering higher participation rates.

Studies incorporating upper body and core exercises showed enhanced outcomes in balance and functional performance. Aqua-cycling provided cardiovascular and musculoskeletal benefits, and was particularly beneficial in obese individuals or those with joint sensitivity.

The psychological impact—including reduced stress and enhanced enjoyment—further supports aquatic therapy's integration into comprehensive OA management plans. Despite methodological variation across studies, the overall direction of evidence is consistent and robust.

Future studies should address protocol standardization, long-term follow-up, and inclusion of broader demographic populations.

## CONCLUSION

Aquatic therapy consistently demonstrates clinical value in managing knee osteoarthritis by alleviating pain, improving physical function, and enhancing quality of life. Its low-impact nature, combined with its adaptability and acceptability, makes it a compelling alternative or adjunct to conventional therapy. Based on current evidence, aquatic therapy should be actively considered in treatment planning for older adults and patients with mobility limitations. Ongoing research is needed to establish long-term outcomes and optimize protocol design.

## REFERENCES

- Dong R, Wu Y, Xu S, Zhang L, Ying J, Jin H, Wang P, Xiao L, Tong P. Is aquatic exercise more effective than land-based exercise for knee osteoarthritis?. *Medicine*. 2018 Dec 1;97(52):e13823.
- Dias JM, Cisneros L, Dias R, Fritsch C, Gomes W, Pereira L, Santos ML, Ferreira PH. Hydrotherapy improves pain and function in older women with knee osteoarthritis: a randomized controlled trial. *Brazilian journal of physical therapy*. 2017 Nov 1;21(6):449-56.
- Rewald S, Lenssen AT, Emans PJ, de Bie RA, van Breukelen G, Mesters I. Aquatic cycling improves knee pain and physical functioning in patients with knee osteoarthritis: a randomized controlled trial. *Archives of physical medicine and rehabilitation*. 2020 Aug 1;101(8):1288-95.
- Kunduracilar Z, Sahin HG, Sonmezer E, Sozay S. The effects of two different water exercise trainings on pain, functional status and balance in patients with knee osteoarthritis. *Complementary therapies in clinical practice*. 2018 May 1;31:374-8.
- Garbi FP, Rocha Júnior PR, Pontes ND, Oliveira AD, Barduzzi GD, Boas PJ. Aquatic physiotherapy in the functional capacity of elderly with knee osteoarthritis. *Fisioterapia em movimento*. 2021 Sep 15;34:e34119.
- Lim JY, Tchai E, Jang SN. Effectiveness of aquatic exercise for obese patients with knee osteoarthritis: a randomized controlled trial. *Pm&r*. 2010 Aug 1;2(8):723-31.
- WYATT FB, MILAM S, MANSKE RC, DEERE R. The effects of aquatic and traditional exercise programs on persons with knee osteoarthritis. *The Journal of Strength & Conditioning Research*. 2001 Aug 1;15(3):337-40.
- Karimi S, Derakhshan M, Tondro A. The effect of aquatherapy on pain intensity and performance in patients with osteoarthritis. *Journal of Family Medicine and Primary Care*. 2024 May 1;13(5):1793-6.
- Noor K, Ahsan MA, Bibi S, Mumtaz T. The Effectiveness of Aquatic Exercises in Improving Balance and Gait in Patients with Knee Osteoarthritis: A Systematic Review. *Pakistan Journal of Medical & Health Sciences*. 2023 Dec 20;17(06):416-.
- Amras A, Kamalakannan R. Role of Aquatic Therapy in Knee Rehabilitation: A Narrative review. *Indian Journal of Physiotherapy and Occupational Therapy*. 2023 Oct 1;17(4).
- Rewald S, Mesters I, Lenssen AF, Emans PJ, Wijnen W, de Bie RA. Effect of aqua-cycling on pain and physical functioning compared with usual care in patients with knee osteoarthritis: study protocol of a randomised controlled trial. *BMC musculoskeletal disorders*. 2016 Dec;17:1-4.
- Wang TJ, Lee SC, Liang SY, Tung HH, Wu SF, Lin YP. Comparing the efficacy of aquatic exercises and land-based exercises for patients with knee osteoarthritis. *Journal of clinical nursing*. 2011 Sep;20(17-18):2609-22.
- Kumar P. Improving IMRaD for writing research articles in social, and health sciences. *International Research Journal of Economics and Management Studies IRJEMS*. 2023;2(1).