



NURTURING MINDS IN EARLY CHILDHOOD: BEHAVIOURAL DISORDERS AMONG TODDLERS AND THE PIVOTAL ROLE OF THE MATERNAL FIGURE

Deepika Verma¹, Dr. Naveen Kumar Jaiswal²

¹PhD Research Scholar, ²Research Supervisor,
Index Nursing College,
Malwanchal University, Indore M.P

ABSTRACT

Early childhood is a critical period for emotional, social, and cognitive development. Behavioural disorders during toddlerhood can significantly influence a child's long-term psychological well-being, academic performance, and social adjustment. This review article explores the prevalence, types, causes, and consequences of behavioural disorders among toddlers, with special emphasis on the role of the maternal figure in prevention, early identification, and management. Drawing from contemporary research, this paper highlights how maternal attachment, mental health, parenting practices, and socio-cultural context shape toddler behaviour. The article also discusses evidence-based interventions and nursing implications. By synthesizing current literature, this review aims to guide healthcare professionals, educators, and caregivers in promoting healthy behavioural development during early childhood.

KEYWORDS: Behavioural disorders, toddlers, maternal role, attachment, parenting practices, early childhood development, mental health, nursing intervention

INTRODUCTION

Toddlerhood, typically defined as the age between one and three years, represents a period of rapid physical, emotional, and cognitive growth. During this stage, children develop autonomy, language skills, emotional regulation, and social awareness. However, this phase is also characterized by vulnerability to behavioural difficulties, as toddlers often struggle to express emotions and adapt to environmental demands. Behavioural disorders during this stage may manifest as excessive aggression, temper tantrums, anxiety, withdrawal, hyperactivity, or oppositional behaviour.

Globally, the prevalence of early childhood behavioural problems is increasing, partly due to changing family structures, urbanization, digital exposure, and psychosocial stressors. These disorders, if left unaddressed, may persist into adolescence and adulthood, leading to academic failure, mental health disorders, and social maladjustment. Therefore, early identification and intervention are essential.

Among various environmental influences, the maternal figure plays a central role in shaping early behavioural patterns. Mothers are often primary caregivers and serve as emotional regulators, attachment figures, and social role models. Their


psychological well-being, parenting style, and responsiveness significantly influence a toddler's behavioural development. This review examines behavioural disorders in toddlers and critically analyzes the role of the maternal figure in fostering positive behavioural outcomes.

Conceptual Framework of Behavioural Disorders in Toddlerhood

Behavioural disorders in toddlers are defined as persistent patterns of disruptive, maladaptive, or emotionally dysregulated behaviours that interfere with daily functioning and social relationships. These behaviours go beyond age-appropriate developmental variations and persist across settings.

Developmental theories provide a foundation for understanding toddler behaviour. Erikson's stage of autonomy versus shame and doubt emphasizes the child's need for independence, while Bowlby's attachment theory highlights the importance of secure caregiver-child relationships. Bandura's social learning theory explains how children imitate observed behaviours. Together, these frameworks suggest that behavioural disorders arise from complex interactions between biological predispositions and environmental influences.

Understanding Behavioural Disorders in Toddlerhood



The biopsychosocial model further explains how genetic vulnerability, neurological development, family environment, and socio-cultural factors interact to influence behaviour. In toddlerhood, limited emotional regulation skills and immature cognitive processes increase susceptibility to behavioural disturbances.

Classification of Behavioural Disorders in Toddlers

Behavioural disorders in toddlers can be broadly classified into externalizing and internalizing disorders. Externalizing disorders involve outwardly directed behaviours, while internalizing disorders involve inward emotional distress.

Table 1: Classification of Behavioural Disorders in Toddlers

Category	Disorder Type	Key Features
Externalizing	Aggressive behaviour	Hitting, biting, temper tantrums
Externalizing	Oppositional behaviour	Defiance, refusal to follow instructions
Externalizing	Hyperactivity	Excessive movement, impulsivity
Internalizing	Anxiety disorders	Fearfulness, separation anxiety
Internalizing	Withdrawal	Social isolation, reduced interaction
Internalizing	Depressive symptoms	Sadness, irritability, loss of interest

Externalizing behaviours are more commonly observed in toddlers due to immature impulse control. Internalizing symptoms are often underdiagnosed, as toddlers may lack verbal skills to express emotional distress.

Gender differences are evident, with boys more likely to display externalizing behaviours and girls more prone to internalizing symptoms. However, environmental factors often mediate these differences.

Epidemiology and Prevalence

Studies indicate that approximately 10–20% of toddlers worldwide exhibit clinically significant behavioural problems. Prevalence varies according to socio-economic status, parental education, family structure, and cultural norms. Children from low-income households and those exposed to parental conflict or neglect demonstrate higher rates of behavioural disorders.

Etiology and Risk Factors

Behavioural disorders in toddlers result from multifactorial influences. Biological factors include genetic predisposition, prenatal exposure to substances, birth complications, and neurodevelopmental delays. Children born preterm or with low birth weight are at higher risk.

In developing countries, limited access to mental health services contributes to underreporting. Cultural perceptions of child behaviour also influence diagnosis. In some societies, aggressive or hyperactive behaviours may be considered normal, delaying professional intervention.

Psychological factors such as temperament, emotional sensitivity, and self-regulation capacity also contribute. Difficult temperament increases vulnerability when combined with inconsistent caregiving.

Environmental factors play a dominant role. Family dysfunction, maternal depression, domestic violence, poverty, and lack of social support adversely affect behavioural



outcomes. Excessive screen time, inadequate stimulation, and disrupted routines further exacerbate problems.

Table 2: Major Risk Factors for Behavioural Disorders

Domain	Risk Factors
Biological	Genetic vulnerability, prematurity, illness
Psychological	Difficult temperament, poor self-regulation
Familial	Maternal depression, marital conflict
Social	Poverty, neglect, lack of stimulation
Environmental	Digital overexposure, unsafe surroundings

The interaction between these factors determines the severity and persistence of behavioural disorders.

Role of Maternal Attachment in Behavioural Development

Attachment refers to the emotional bond between a child and primary caregiver. Secure attachment develops when mothers respond consistently and sensitively to their child’s needs. Securely attached toddlers demonstrate better emotional regulation, social competence, and stress management.

Insecure attachment patterns, including avoidant, ambivalent, and disorganized attachment, are associated with behavioural disturbances. Children with insecure attachment often exhibit aggression, anxiety, and social withdrawal. Maternal insensitivity, neglect, or inconsistency disrupts attachment formation.

Maternal responsiveness during infancy lays the foundation for later behaviour. Physical affection, verbal interaction, and emotional availability promote trust and self-confidence. Conversely, emotionally unavailable mothers may unintentionally foster insecurity and behavioural instability.

Maternal Mental Health and Toddler Behaviour

Maternal mental health is a critical determinant of child development. Depression, anxiety, and stress impair maternal sensitivity and responsiveness. Depressed mothers may display reduced emotional engagement, irritability, and inconsistent discipline.

Postpartum depression has been strongly linked to behavioural problems in toddlers, including hyperactivity, aggression, and emotional dysregulation. Chronic maternal stress also affects parenting quality and increases family conflict.

Table 3: Impact of Maternal Mental Health on Toddler Behaviour

Maternal Condition	Impact on Parenting	Behavioural Outcome in Toddler
Depression	Low responsiveness	Withdrawal, irritability
Anxiety	Overprotection	Fearfulness, dependency
Chronic stress	Inconsistent discipline	Aggression, impulsivity
Substance abuse	Neglectful caregiving	Emotional instability

Early screening and treatment of maternal mental disorders are essential for promoting healthy child development.

Parenting Practices and Behavioural Regulation

Parenting style significantly influences toddler behaviour. Authoritative parenting, characterized by warmth and consistent discipline, promotes self-control and social competence. Authoritarian parenting, involving strict control and limited emotional support, may lead to fear and rebellion. Permissive parenting often results in poor impulse control and defiance.

Positive parenting techniques such as praise, active listening, and structured routines enhance behavioural regulation. Conversely, harsh punishment, verbal abuse, and neglect increase the risk of behavioural disorders.

Maternal modeling of emotional regulation and problem-solving skills helps toddlers develop adaptive coping strategies. Mothers who demonstrate calm responses to stress foster emotional resilience in children.

Socio-Cultural and Environmental Influences

Maternal caregiving practices are shaped by cultural beliefs, social norms, and economic conditions. In collectivist cultures, extended family support may buffer maternal stress, whereas in nuclear families, mothers often experience caregiving burden.

Urbanization, maternal employment, and reduced family support systems influence child-rearing practices. Working mothers may face challenges in balancing professional and caregiving roles, leading to emotional exhaustion.

Digital media exposure has emerged as a significant environmental factor. Excessive screen time reduces parent-child interaction and contributes to attention problems and emotional dysregulation.

Assessment and Early Identification

Early identification of behavioural disorders requires systematic screening and observation. Healthcare professionals use standardized tools such as the Child Behavior Checklist and Ages and Stages Questionnaires.

Maternal reports, clinical interviews, and developmental assessments provide comprehensive evaluation. Early warning signs include persistent aggression, lack of eye contact, sleep disturbances, feeding problems, and social withdrawal.

Nurses and primary care providers play a crucial role in monitoring developmental milestones and guiding mothers regarding behavioural concerns.

Intervention Strategies and Maternal Involvement


Intervention for toddler behavioural disorders should be family-centered and developmentally appropriate. Parent training programs enhance maternal skills in behaviour management, communication, and emotional support.

Cognitive-behavioural approaches adapted for young children focus on emotional awareness and self-regulation. Play therapy

and parent-child interaction therapy strengthen attachment and communication.

Community-based programs provide social support and mental health services to vulnerable mothers. Home visiting programs have demonstrated positive outcomes in improving parenting practices and child behaviour.

Unveiling Intervention Strategies for Toddler Behavioural Disorders



Maternal empowerment through education and counseling is essential for sustainable behavioural improvement.

Nursing Implications and Professional Role

Nurses play a vital role in promoting maternal and child mental health. They conduct developmental screenings, provide parenting education, and offer emotional support to mothers.

Community health nurses facilitate parent support groups and coordinate referrals to mental health professionals. In hospital and primary care settings, nurses advocate for early intervention services.

Integrating mental health services into maternal and child healthcare programs enhances accessibility and reduces stigma.

Challenges and Future Directions

Despite growing awareness, several challenges remain. Limited mental health resources, cultural stigma, and lack of trained professionals hinder early intervention. In low-resource settings, behavioural problems are often neglected.

Future research should focus on longitudinal studies, culturally sensitive interventions, and digital health solutions.

Strengthening maternal mental health services and parenting support programs is essential.

Interdisciplinary collaboration among healthcare providers, educators, and policymakers is necessary to address behavioural disorders comprehensively.

CONCLUSION

Behavioural disorders among toddlers represent a significant public health concern with long-term implications. The maternal figure plays a pivotal role in shaping behavioural development through attachment, mental health, and parenting practices. Secure attachment, positive parenting, and maternal psychological well-being serve as protective factors, while stress, depression, and inconsistent caregiving increase vulnerability.

Early identification, family-centered interventions, and strong nursing support systems are crucial for preventing and managing behavioural disorders. Empowering mothers through education, counseling, and social support can foster healthy emotional and behavioural development in toddlers. Addressing maternal and child mental health holistically will contribute to nurturing resilient and well-adjusted future generations.



REFERENCES

1. Bowlby J. *Attachment and Loss. Vol. 1. New York: Basic Books; 1982.*
2. Erikson EH. *Childhood and Society. New York: Norton; 1993.*
3. Ainsworth MDS, Blehar MC, Waters E, Wall S. *Patterns of Attachment. Hillsdale: Erlbaum; 1978.*
4. Sheldrick RC, Marakovitz S, Garfinkel D, Perrin EC, Carter AS. *Prevalence and co-occurrence of developmental and emotional-behavioral problems in young children. Academic pediatrics. 2023 Apr 1;23(3):623-30.*
5. Goodman SH, Rouse MH, Connell AM, et al. *Maternal depression and child psychopathology. Psychol Bull. 2011;137(1):124-155.*
6. Bacherini A, Iglizzo R, Cagiano R, Mancini A, Tancredi R, Muratori F, Balboni G. *Behavioral and emotional problems of toddlers with autism spectrum disorder: Effects of parents' sociocultural level and individual factors. Research in Developmental Disabilities. 2021 Dec 1;119:104106.*
7. Campbell SB. *Behavior problems in preschool children. Dev Psychopathol. 2002;14(3):467-495.*
8. Feldman R. *Parent-infant synchrony. Curr Dir Psychol Sci. 2007;16(6):340-345.*
9. Sanders MR. *Triple P-Positive Parenting Program. Clin Child Fam Psychol Rev. 1999;2(2):71-90.*
10. Shonkoff JP, Phillips DA. *From Neurons to Neighborhoods. Washington DC: National Academy Press; 2000.*
11. Sadeghi S, Pouretamad HR, Badv RS, Brand S. *Associations between symptom severity of autism spectrum disorder and screen time among toddlers aged 16 to 36 months. Behavioral Sciences. 2023 Feb 27;13(3):208.*
12. WHO. *Improving Early Childhood Development. Geneva: WHO; 2020.*
13. Landry SH, Smith KE, Swank PR. *Responsive parenting. Dev Psychol. 2006;42(4):627-642.*