



UNRAVELLING DYSMENORRHEA: A REVIEW OF CURRENT EVIDENCE AND PRACTICE IMPLICATIONS IN PHARMACY PRACTICE

Alwy Jose, Malavika P.U, Ayisha K.M

Article DOI: <https://doi.org/10.36713/epra27655>

DOI No: 10.36713/epra27655

ABSTRACT

Primary dysmenorrhea remains one of the most prevalent gynaecological disorders affecting adolescents and young women globally. Contemporary cross-sectional and analytical studies from Asia, the Middle East, Europe, and South America consistently report high prevalence rates with significant academic absenteeism, reduced work productivity, impaired quality of life, and economic burden. Pathophysiological advances highlight the central role of prostaglandin-mediated uterine hypercontractility and inflammatory cascades, with emerging insights into immune and stress-related mechanisms. Validated tools such as the Working ability, Location, Intensity, Days of pain, Dysmenorrhea (WaLIDD) score demonstrate strong reliability and cross-cultural applicability, facilitating standardized severity assessment. Non-steroidal anti-inflammatory drugs remain first-line pharmacotherapy; however, variations in utilization patterns, self-medication practices, and safety concerns persist. Network meta-analyses confirm NSAID superiority, while evidence for exercise, stretching, and complementary therapies is growing yet heterogeneous. Psychological stress, nutritional status, body mass index, and lifestyle factors consistently emerge as significant modifiers of symptom severity. Despite expanding literature, gaps remain in longitudinal data, comparative effectiveness research, and integration of patient-reported outcomes into routine care. A multidimensional, evidence-based framework that integrates pharmacologic therapy, lifestyle modification, and psychosocial support is essential for optimized real-world management.

KEYWORDS: Primary dysmenorrhea; WaLIDD score; NSAIDs; Quality of life; non-pharmacological therapy

INTRODUCTION

Background

Primary dysmenorrhea is defined as painful menstruation in the absence of identifiable pelvic pathology. Epidemiological studies across medical and non-medical student populations demonstrate prevalence rates frequently exceeding 60–80% [18,41,57]. The burden extends beyond pain, encompassing absenteeism, reduced productivity, psychological distress, and diminished quality of life [9,13,27].

Despite its high prevalence, dysmenorrhea remains underprioritized in clinical and public health frameworks. Increasing evidence demonstrates its association with stress [12,17], socioeconomic disparities [3,24], lifestyle factors [38], and nutritional determinants [28].

Scope

This review integrates findings from observational studies, validation research, systematic reviews, and network meta-analyses published between 2015 and 2025, including validation of severity scales such as the WaLIDD score [49] and therapeutic comparisons [50].

Rationale and Need

The heterogeneity of management practices, widespread self-medication [52,56], and growing evidence supporting multimodal therapy mandate a consolidated, evidence-based perspective bridging severity assessment and real-world management.

OBJECTIVES

1. To summarize global prevalence and determinants of primary dysmenorrhea.
2. To evaluate validated severity assessment tools including WaLIDD and Numerical Rating Scale.
3. To compare pharmacological and non-pharmacological management strategies.
4. To assess quality of life, productivity, and economic implications.
5. To identify controversies, methodological limitations, and research gaps.

METHODOLOGY OF LITERATURE SEARCH

A structured literature search was conducted across PubMed, Scopus, Google Scholar, and Web of Science databases covering publications from 2015 to 2025, with selective inclusion of landmark earlier studies. Keywords included “primary dysmenorrhea,” “WaLIDD score,” “NSAIDs,” “quality of life,” “exercise therapy,” and “economic burden.” Cross-sectional studies, systematic reviews, meta-analyses, validation studies, and drug utilization studies published in English were included. Studies focusing on secondary dysmenorrhea, case reports, and non-peer-reviewed commentaries were excluded unless methodologically significant. This approach ensured comprehensive coverage of epidemiology, assessment tools, therapeutic interventions, and real-world management patterns.



DISCUSSION

Epidemiology and Determinants

Recent cross-sectional surveys across India, Indonesia, Pakistan, Malaysia, Greece, and Saudi Arabia report persistent high prevalence [1,18,30,45]. BMI, physical inactivity, stress, and poor nutritional status correlate significantly with severity [11,12,28,38]. War displacement and mental health vulnerability amplify severity in conflict settings [24,32].

Pathophysiology and Emerging Mechanisms

Traditional understanding implicates excess endometrial prostaglandins causing uterine hypercontractility [35]. Novel inflammatory biomarkers identified in menstrual effluent expand the mechanistic model toward immune modulation [7]. Psychoneuroendocrine pathways linking stress and pain sensitization are increasingly recognized [12,17].

Diagnostic Evaluation and Differential Diagnosis

Although primary dysmenorrhea is diagnosed clinically, exclusion of secondary causes such as endometriosis, adenomyosis, pelvic inflammatory disease, and uterine fibroids remains essential in refractory or atypical presentations. Red flag indicators include progressive pain, dyspareunia, abnormal uterine bleeding, and poor response to NSAIDs. Clinical history, pelvic examination (when appropriate), and ultrasonography are recommended for diagnostic clarification. Early differentiation optimizes therapeutic outcomes and prevents delayed identification of underlying pathology.

Severity Assessment Tools

The WaLIDD score, introduced by Teherán et al. [49], integrates multidimensional parameters. Cross-cultural validation studies in Arabic, Turkish, and Brazilian populations confirm reliability and validity [14,25,26]. Comparative studies demonstrate concordance between WaLIDD and Numerical Rating Scale but superior multidimensional profiling with WaLIDD [39].

Pharmacological Management

NSAIDs remain first-line therapy by inhibiting cyclooxygenase-mediated prostaglandin synthesis. Network meta-analysis confirms superior efficacy over placebo [50]. However, drug utilization studies reveal irrational patterns, delayed initiation, and inappropriate dosing [23,48]. Self-medication remains widespread among students [52,56], raising safety concerns.

Complementary therapies, including homoeopathy [21] and alternative approaches [44], demonstrate variable evidence quality. Hormonal therapy and combined approaches are underreported in many regional studies.

Non-Pharmacological Interventions

Exercise and stretching significantly reduce severity scores in randomized and quasi-experimental designs [2,33]. A 2025 network meta-analysis supports the effectiveness of non-pharmacological interventions, though heterogeneity limits definitive ranking [6]. Lifestyle modification demonstrates protective effects [8,38].

Quality of Life and Economic Burden

Quality of life impairment is consistently documented [13,27]. Financial consequences include direct treatment costs and productivity loss [19,53]. Academic absenteeism remains a critical metric among students [41].

Clinical Implications for Gynaecological Practice

Translation of evidence into routine practice requires standardized pain assessment, rational NSAID prescribing, early initiation at onset of menses, and patient counseling regarding dosing adherence. Incorporating validated tools such as the WaLIDD score into outpatient settings enables objective monitoring. Multimodal management—including pharmacotherapy, structured exercise, stress reduction, and lifestyle modification—should be positioned as the default care pathway rather than sequential trial-and-error management.

Public Health and Policy Implications

Given its high prevalence and documented impact on productivity and educational attendance, dysmenorrhea warrants recognition as a public health priority. Institutional policies addressing menstrual health education, workplace flexibility, and affordable access to analgesics are justified. Economic evaluations indicate measurable productivity loss, reinforcing the need for policy-level intervention and inclusion of menstrual health within reproductive health programs.

Strengths and Limitations of Current Evidence

The reviewed literature provides robust prevalence data and consistent associations with lifestyle and psychological variables. However, the predominance of cross-sectional designs limits causal inference. Variability in outcome measures and underreporting of adverse drug reactions reduce external validity. Limited randomized controlled trials comparing pharmacological and non-pharmacological interventions restrict definitive clinical hierarchy establishment.

Trends, Controversies and Limitations

The literature is dominated by cross-sectional designs, limiting causal inference [55]. Heterogeneity in severity scales reduces comparability. Many studies lack long-term follow-up and pharmacoeconomic evaluation. Complementary therapies remain controversial due to methodological limitations [21,47].

Research Gaps

- Need for longitudinal cohort studies.
- Standardization of severity assessment tools globally.
- Comparative effectiveness research between NSAIDs and structured exercise programs.
- Integration of patient-reported outcomes into routine gynaecological practice.

Future Research Directions

Future investigations should prioritize longitudinal cohort studies to establish temporal associations and risk prediction models. Standardization of severity scales across global populations is necessary to improve comparability. Comparative effectiveness trials evaluating NSAIDs versus structured exercise protocols, and integrated multimodal



interventions, are warranted. Additionally, pharmacoeconomic analyses and incorporation of patient-reported outcome measures should become standard endpoints in dysmenorrhea research.

CONCLUSION

Primary dysmenorrhea is a high-burden, multidimensional disorder with substantial academic, occupational, psychological, and economic implications. Evidence consistently supports prostaglandin-mediated pathophysiology with emerging inflammatory and stress-related mechanisms. The WaLIDD score represents a validated, cross-culturally adaptable severity assessment tool that enhances multidimensional evaluation. NSAIDs remain the cornerstone of therapy; however, irrational use and self-medication persist. Non-pharmacological interventions, particularly structured exercise, demonstrate meaningful benefit and should be integrated into management algorithms. Current literature is constrained by cross-sectional methodologies and heterogeneity in outcome measures. Future research must prioritize longitudinal designs, standardized assessment frameworks, and real-world comparative effectiveness studies. A comprehensive, patient-centered, multimodal strategy is imperative to transition dysmenorrhea management from symptomatic relief toward optimized, evidence-driven care pathways.

ACKNOWLEDGEMENT

I express sincere gratitude to **Dr. S. Aravinth Vijay Jesuraj**, my research guide, and all faculty members of the Department of Pharmacy Practice, Nirmala College of Health Science, for their academic guidance and continuous support.

REFERENCES

- Valiyaparambil JA, More PD, Bahekar SE. Menstrual health challenges: A rural cross-sectional study on dysmenorrhea among medical students of Maharashtra. *Al Ameen J Med Sci*. 2025 Jul 1;18(3):192-7.
- Rajan Balakrishnan, Nelson Arputharaj and Lai Wen Qui. Effect of stretching exercise on primary dysmenorrhea among young female adults. *Int J Phys Educ Sports Health*. 2025 Sep 21;12(6):324-7. doi:<https://www.doi.org/10.22271/kheljournal.2025.v12.i6e.4109>
- Panigrahi J, Rout D, Jena B, Dash P, Mohanty NR. Dysmenorrhea decoded: Unravelling the complex interplay of pathophysiology, socioeconomic disparities and therapeutic innovations. *Int J Appl Res*. 2025 Jan 1;11(1):182-7. doi:[10.22271/allresearch.2025.v11.i1c.12275](https://doi.org/10.22271/allresearch.2025.v11.i1c.12275)
- Nurwany R, Ramadhani NT, Putri SSF, Zulissetiana EF. Physical Activity Impact on Severity of Dysmenorrhea among Junior High School Students: A Cross Sectional Study. *Obstet Gynecol*. 2025;13(2).
- Magar NT, Regmi S, Adhikari A, Dev B, Timsina J, Das SB. Assessment and Comparison on Use of Non-Steroidal Anti-Inflammatory Drugs for the Management of Dysmenorrhea by Medical and Non-Medical Students. *J Drug Delivery Ther*. 2025 Apr 15;15(4):80-4. doi:[10.22270/jddt.v15i4.7074](https://doi.org/10.22270/jddt.v15i4.7074)
- Liu J, Wang Y, Zhang J, Fan X, Chen H, Zuo G, et al. Efficacy and Safety of Non-Pharmacological Therapies for Primary Dysmenorrhea: A Network Meta-Analysis. *JPR*. 2025 Feb;Volume 18:975-91. doi:[10.2147/JPR.S498184](https://doi.org/10.2147/JPR.S498184)
- Kyathanahalli CN, Tu F, Hellman K. Inflammatory Mechanisms of Dysmenorrhea: Novel Insights from Menstrual Effluent in an Adolescent Cohort [Internet]. Preprints; 2025 [cited 2026 Jan 22]. (132). Available from: <https://doi.org/10.1111/1471-0528.18275> doi:[10.22541/au.174167500.02946487/v1](https://doi.org/10.22541/au.174167500.02946487/v1)
- Kumuda T. The Impact of Lifestyle Factors on Menstrual Irregularities in Adolescent Girls: A Prospective Observational Study. *Journal of Contemporary Clinical Practice*. 2025 Aug 9;11(9):124-31. doi:[10.61336/jccp/25-09-18](https://doi.org/10.61336/jccp/25-09-18)
- Kour G, Katoch S. Prevalence and impact of dysmenorrhea on quality of life among women: a cross-sectional study. *International Journal of Medicine and Public Health*. 2025;15(3):2017-21. doi:[10.70034/ijmedph.2025.3.373](https://doi.org/10.70034/ijmedph.2025.3.373)
- Keerthana M, Kumar JP, Charitha PS, Deepthi ELP, Bhavana M, Anusha A. Manifestation and Management of Dysmenorrhea - A Cross Sectional Study. *IJST*. 2025 Jun 17;18(22):1782-91. doi:[10.17485/IJST/v18i22.482](https://doi.org/10.17485/IJST/v18i22.482)
- Fatima N, Saeed A, Khurshid N, Shahzadi N, Rafi MA. Frequency of primary dysmenorrhea among young females and its association with body mass index and physical activity: A cross-sectional study from Islamabad and Rawalpindi, Pakistan. *Rawal Medical Journal*. 2025 Sep;50(3):644-7.
- De Moraes PC, Avila MA, Firão CB, De Sousa VPS, Driusso P. Perceived stress is associated with primary dysmenorrhea in Brazilian women: a cross-sectional study. *BMC Public Health*. 2025 Apr 5;25(1):1290 (9 pages). doi:[10.1186/s12889-025-21804-6](https://doi.org/10.1186/s12889-025-21804-6)
- Dar M, Khan A, Shah SS, Aleem A, Jaber AAS, Hussain M, et al. Evaluation of health-related quality of life of female students suffering from primary dysmenorrhea: findings of a cross-sectional study from Pakistan. *Front Public Health*. 2025 Apr 24;13:1467377. doi:[10.3389/fpubh.2025.1467377](https://doi.org/10.3389/fpubh.2025.1467377)
- Arikan H, Erol E. Working ability, Location, Intensity, Days of Pain, Dysmenorrhea (WaLIDD): cross-cultural adaptation, reliability, and validity of the Turkish version. *BMC Women's Health*. 2025 Mar 22;25(1):133. doi:[10.1186/s12905-025-03673-8](https://doi.org/10.1186/s12905-025-03673-8)
- Ankita Singh, Rashmi Kumari, Smita Bharti, Neha Kumari, Sveta Kumari, Annu Priya, Luckey Kumari, Sneha Singh, Mona Singh, Anushreya Soreng, Pinki Kumari, Archana Kumari, Dibya Kumari, Priyanka Kumari, Sneha Sinha,, Jasmine Kumari, Chhaya Kumari, Pushpa Kumari, Arnab Roy*. Unravelling the Complexity of Dysmenorrhea: A Review of the Underlying Mechanisms and Treatment Options for Caregivers and Patients. *International journal of scientific research and technology [Internet]*. 2025;2(10):245-52. Available from: <https://doi.org/10.5281/zenodo.17339137>
- Alali Z. From Pain to Impairment: A Study of the Prevalence, Severity, and Impact of Dysmenorrhea in Women From the Eastern Region of Saudi Arabia. *Clin Exp Obstet Gynecol*. 2025 Nov 18;52(11):41538. doi:[10.31083/CEOG41538](https://doi.org/10.31083/CEOG41538)
- Abdulkhalik MS, Alshahrani MA, Alamri GMA, Aldawodi RM, Memish HA, Asiri SA, et al. The Impact of Stress and Anxiety on Dysmenorrhea Severity: A systematic review. *SMHJ*. 2025 Dec 9;6(1):156-65. doi:[10.54293/smhj.v6i1.185](https://doi.org/10.54293/smhj.v6i1.185)



18. Situmorang H, Sutanto RL, Tjoa K, Rivaldo R, Adrian M. Prevalence and risk factors of primary dysmenorrhoea among medical students: a cross-sectional survey in Indonesia. *BMJ Open*. 2024 Oct;14(10):e086052. doi:10.1136/bmjopen-2024-086052
19. Santhoshkumar S. G. The Financial Ramifications of Dysmenorrhea: A Cross-Sectional Study on Employed Women in Chennai. *International Journal of Innovative Scientific Research*. 2024;2(4):15–20. doi:105281/ijisr550724
20. Mody V, Patel R, Patel H, Dave J, Vasava P. Primary dysmenorrhea and its association with psychological symptoms among college going adolescent girls - a cross sectional. *International Journal of Current Science (IJCSPUB)*. 2024;14(2). doi:IJCSP24B1002
21. Lata S. Efficacy Of Homoeopathic Medicines On Dysmenorrhoea. *JRTDD*. 2024 Sep 20;6(7):1322–6. doi:10.53555/jrtdd.v6i7s.3376
22. Kirsch E, Rahman S, Kerolus K, Hasan R, Kowalska D, Desai A, et al. Dysmenorrhea, a Narrative Review of Therapeutic Options. *JPR*. 2024 Aug;Volume 17:2657–66. doi:10.2147/JPR.S459584
23. Gandhi S, Dawane JS. Study the pattern of drug use in dysmenorrhea, its effectiveness and problems faced during therapies in young female. *Journal of Chemical Health Risks*. 2024 Jun 3;14(2):2591–7.
24. Gammoh O, Al Rob OA, Alqudah A, Al-Smadi A, Dobain MO, Zeghoul R, et al. Risk factors for severe dysmenorrhea in Arab women: A focus on war displacement and mental health outcomes. *AIMSPH*. 2024;11(1):209–22. doi:10.3934/publichealth.2024010
25. Bou Haidar P, Alam D, Obeid S, Malaeb D, Dabbous M, Sakr F, et al. Validation of the Working Ability, Location, Intensity, Days of Pain and Dysmenorrhea (VWaLIDD) Scale in Arabic-Speaking Young Female Adults. *IJWH*. 2024 Sep 27;Volume 16:1565–71. doi:10.2147/IJWH.S468803
26. Avila MA, Arruda GTD, Godoy AGD, Driusso P. Translation, cross-cultural adaptation to Brazilian Portuguese and measurement properties of the WaLIDD score. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics*. 2024 Mar 14;46:e-rbgo16. doi:10.61622/rbgo/2024AO16
27. Amza M, Findekle S, Haj Hamoud B, Sima RM, Poenaru MO, Popescu M, et al. Dysmenorrhea and Its Impact on Patients' Quality of Life – A Cross-Sectional Study. *JCM*. 2024 Sep 24;13(19):5660. doi:10.3390/jcm13195660
28. Ronny, Nur Alvoionita Tianti Kusuma, Vidi Posdo A. Simarmata. Correlation between nutritional status and primary dysmenorrhea based on work ability, location, intensity, days of pain, and dysmenorrhea (Walidd) Score. *World J Bio Pharm Health Sci*. 2023 May 30;14(2):046–62. doi:10.30574/wjbphs.2023.14.2.0184
29. Omar Salem G. Dysmenorrhea severity in war refugees with hypertension: a cross-talk with antihypertensives and analgesics. *Afr H Sci*. 2023 Dec 27;23(4):432–7. doi:10.4314/ahs.v23i4.47
30. Mohamad Bakro R, Farrukh MJ, Rajagopal M, Kristina SA, Ramatillah DL, Ming LC, et al. Assessment of prevalence, knowledge and health-related practices of dysmenorrhea among Malaysian women in Kuala Lumpur: a cross-sectional survey. *Annals of Medicine*. 2023 Dec 12;55(2):2281655. doi:10.1080/07853890.2023.2281655
31. Juwitasari, Dhea Erlita R, Harini R. Assessment of Dysmenorrhea using WaLIDD Score Instruments on Nursing Students. *Kme*. 2023 Jun 23;145–51. doi:10.18502/kme.v3i3.13498
32. Ghandour R, Hammoudeh W, Stigum H, Giacaman R, Fjeld H, Holmboe-Ottesen G. Menstrual characteristics and dysmenorrhea among Palestinian adolescent refugee camp dwellers in the West Bank and Jordan: a cross-sectional study [Internet]. In Review; 2023 [cited 2026 Feb 23]. Available from: <https://www.researchsquare.com/article/rs-2415553/v1> doi:10.21203/rs.3.rs-2415553/v1
33. Anis T, Hasan S, Lohana N, Salahuddin I, Saeed H. Effects of Aerobic Program on the Severity of Dysmenorrhea using WaLIDD Score. *Journal of Xi'an Shiyou University, Natural Science Edition* [Internet]. 2023 Jun 6;19(6):550–4. Available from: <http://xisdxjxsu.asia>
34. Parekh S. Prevalence of dysmenorrhea and associated symptoms among adolescent girls: a cross sectional study. *Journal of Emerging Technologies and Innovative Research (JETIR)*. 2022;9(3):b668–96. doi:JETIR2203186
35. Itani R, Soubra L, Karout S, Rahme D, Karout L, Khojah HMJ. Primary Dysmenorrhea: Pathophysiology, Diagnosis, and Treatment Updates. *Korean J Fam Med*. 2022 Mar 20;43(2):101–8. doi:10.4082/kjfm.21.0103
36. De Arruda GT, Driusso P, Rodrigues JC, de Godoy AG, Avila MA. Numerical rating scale for dysmenorrhea-related pain: a clinimetric study. *Gynecological Endocrinology*. 2022 Aug 3;38(8):661–5. doi:10.1080/09513590.2022.2099831
37. Algabbani AM, Alzahrani SA, Almomen SM, Hafiz RA. Readability of information imprinted in patient information leaflets (PILs) in Saudi Arabia: The case of antihypertensive medications. *Exploratory Research in Clinical and Social Pharmacy*. 2022 Dec;8:100179. doi:10.1016/j.rcsop.2022.100179
38. Al-Husban N, Odeh O, Dabit T, Masadeh A. The Influence of Lifestyle Variables on Primary Dysmenorrhea: A Cross-Sectional Study. *IJWH*. 2022 Apr;Volume 14:545–53. doi:10.2147/IJWH.S338651
39. Rashidi Fakari F, Simbar M, Tahmasebi G, Ebadi A, Rashidi Fakari F, Nasiri M, et al. Efficacy of Working Ability, Location, Intensity, Days of Pain, Dysmenorrhea (WaLIDD) and Verbal Rating Scale (Pain and Drug) in Diagnosing and Predicting Severity of Dysmenorrhea among Adolescents: A Comparative Study. *J Obstet Gynecol Cancer Res*. 2021 May 5;6(2):81–6. doi:10.30699/jogcr.6.2.81
40. Parisa N, M. Khairul Kahfi Pasaribu, Bintang Arroyantri Prananjaya, Ardesy Melizah Kurniati, Ziske Maritska. Patterns of Using Analgesics for Menstrual Pain in Female Students Sriwijaya University Faculty of Medicine. *AMCR*. 2021 Oct 13;1(2):49–57. doi:10.37275/amcr.v1i2.10
41. Syed A, Rao SB. Prevalence of premenstrual syndrome and dysmenorrhea among medical students and its impact on their college absenteeism. *Int J Reprod Contracept Obstet Gynecol*. 2020 Mar 25;9(4):1377. doi:10.18203/2320-1770.ijrcog20201042
42. Priya Sharma, Hetal Patel*, Sujeeta Malik. Descriptive study to assess the practice of women on management of dysmenorrhea in selected areas of d&nh. *Global journal for research analysis*. 2020 Jul;9(7):61–3. doi:10.36106/gjra
43. Kemigisha E, Rai M, Mlahagwa W, Nyakato VN, Ivanova O. A Qualitative Study Exploring Menstruation



- Experiences and Practices among Adolescent Girls Living in the Nakivale Refugee Settlement, Uganda. *IJERPH*. 2020 Sep 11;17(18):6613. doi:10.3390/ijerph17186613
44. Abubakar U, Zulkarnain AI, Samri F, Hisham SR, Alias A, Ishak M, et al. Use of complementary and alternative therapies for the treatment of dysmenorrhea among undergraduate pharmacy students in Malaysia: a cross sectional study. *BMC Complement Med Ther*. 2020 Dec;20(1):285. doi:10.1186/s12906-020-03082-4
45. Vlachou E, Owens DA, Lavdaniti M, Kalemikerakis J, Evangelou E, Margari N, et al. Prevalence, Wellbeing, and Symptoms of Dysmenorrhea among University Nursing Students in Greece. *Diseases*. 2019 Jan 8;7(1):5. doi:10.3390/diseases7010005
46. Sulistyowati DWW, Harumi AM. Determinants of Dysmenorrhea Pain Levels of in Adolescents. *AIJHA*. 2019 Dec 30;3(1):1-13. doi:10.33846/aijha30101
47. Sharghi M, Mansurkhani SM, Ashtary-Larky D, Kooti W, Niksefat M, Firoozbakht M, et al. An update and systematic review on the treatment of primary dysmenorrhea. *JBRA Assisted Reproduction*. 2019;23(1):51-7. doi:10.5935/1518-0557.20180083
48. Nandeshwar MB, Chakravorty AD. A Study of drug utilization pattern in gynaecology out-patient department of a tertiary care hospital. *Int J Res Med Sci*. 2019 Oct 24;7(11):4280. doi:10.18203/2320-6012.ijrms20195002
49. Teherán A, Píneros LG, Pulido F, Mejía Guatibonza MC. WaLIDD score, a new tool to diagnose dysmenorrhea and predict medical leave in university students. *IJWH*. 2018 Jan; Volume 10:35-45. doi:10.2147/IJWH.S143510
50. Feng X, Wang X. Comparison of the efficacy and safety of non-steroidal anti-inflammatory drugs for patients with primary dysmenorrhea: A network meta-analysis. *Mol Pain*. 2018 Jan;14:1-14. doi:10.1177/1744806918770320
51. Mathur R, editor. National ethical guidelines for biomedical and health research involving human participants. New Delhi: Indian Council of Medical Research; 2017.
52. Fatima A, R M, Ambika B, Rajarathna K. Self-medication practice in primary dysmenorrhea among medical and paramedical students - A cross-sectional questionnaire study. *Natl J Physiol Pharm Pharmacol*. 2017;7(4):458-63. doi:10.5455/njppp.2017.7.1233424122016
53. Akiyama S, Tanaka E, Cristeau O, Onishi Y, Osuga Y. Evaluation of the treatment patterns and economic burden of dysmenorrhea in Japanese women, using a claims database. *CEOR*. 2017 May; Volume 9:295-306. doi:10.2147/CEOR.S127760
54. Akiyama S, Tanaka E, Cristeau O, Onishi Y, Osuga Y. Evaluation of the treatment patterns and economic burden of dysmenorrhea in Japanese women, using a claims database. *CEOR*. 2017 May; Volume 9:295-306. doi:10.2147/CEOR.S127760
55. Setia M. Methodology series module 3: Cross-sectional studies. *Indian J Dermatol*. 2016;61(3):261. doi:10.4103/0019-5154.182410
56. B. J, V. A. Comparison of self-medication practice for dysmenorrhoea in medical, nursing and dental students. *Int J Basic Clin Pharmacol*. 2016;269-73. doi:10.18203/2319-2003.ijbcp20160649
57. Kural M, Noor N, Pandit D, Joshi T, Patil A. Menstrual characteristics and prevalence of dysmenorrhea in college going girls. *J Family Med Prim Care*. 2015;4(3):426-31. doi:10.4103/2249-4863.161345
58. Kaur S, Sheoran P, Sarin J. Assessment And Comparison of Dysmenorrhea in Terms of Severity of Pain and Utilization of Non Steroid Anti-Inflammatory Drugs among Unmarried and Married Women. *International Journal of Caring Sciences*. 2015 Dec;8(3):737-46.